arXiv:1405.1900v1 [math.RT] 8 May 2014

FACTORIZATION OF GROUP DETERMINANT IN SOME GROUP ALGEBRAS

NAOYA YAMAGUCHI

ABSTRACT. We give an analog of Frobenius' theorem about the factorization of the group determinant on the group algebra of finite abelian groups and we extend it into dihedral groups and generalized quaternion groups. Furthermore, we describe the group determinant of dihedral groups and generalized quaternion groups as a circulant determinant of homogeneous polynomials.

This analog on the group algebra is stronger than Frobenius's theorem and as a corollary, we obtain a simple expression formula for inverse elements in the group algebra. Furthermore, the commutators of irreducible factors of the factorization of the group determinant on the group algebra corresponding to degree one representations have interesting algebraic properties. From this result, we know that degree one representations form natural pairing.

At the current stage, the extension of Frobeinus' theorem is not represent as a determinant. We expect to find a determinant expression similar to Frobenius' theorem.

1. INTRODUCTION

In this paper, we give an analog of the factorization of the group determinant on the group algebra, where the group G is a finite abelian group, a dihedral group or a generalized quaternion group. The group determinant $\Theta(G)$ is the determinant of the matrix whose elements are independent variables x_g corresponding to $g \in G$. Frobenius gave the following theorem about the factorization of the group determinant.

Theorem 1 (Frobenius)**.** *Let G be a finite group and $\widehat{G}$ be a complete set of irreducible representations of G. Then, we have*

$$\Theta(G) = \prod_{\varphi \in \widehat{G}} \det \left(\sum_{g \in G} \varphi(g) x_g \right)^{\deg \varphi}.$$

Our main results of this paper are non-trivial extensions of this theorem for some group on the corresponding group algebras.

1.1. Results for finite abelian groups.

Our main result for finite abelian group is the following theorem.

Date: May 9, 2014.

2010 *Mathematics Subject Classification.* Primary 20C15; Secondary 15A15; 22D20.

Key words and phrases. group determinant; group algebra; dihedral group; generalized quaternion group; circulant determinant.

Theorem 2. *Let G be a finite abelian group and e be the unit element of G. Then, we have*

$$\Theta(G)e = \prod_{\chi\in\widehat{G}} \sum_{g\in G} \chi(g)x_g g.$$

Let

$$\mathbb{C}G = \left\{ \sum_{g\in G} x_g g \mid x_g \in \mathbb{C}, g \in G \right\}$$

be the group algebra of G. Note that the equality in Theorem 2 is the equality on the group algebra. Theorem 2 is stronger than Theorem 1. In fact, let $F : \mathbb{C}G \to \mathbb{C}$ be a $\mathbb{C}$-linear ring homomorphism such that $F(g) = 1$ for all $g \in G$. Theorem 1 for finite abelian groups follows by applying F to Theorem 2. Moreover, we obtain the following formula for inverse elements in the group algebra.

Corollary 3. *Let G be a finite abelian group and χ_1 be the trivial representation of G. If $\Theta(G) \neq 0$, then we have*

$$\left(\sum_{g\in G} x_g g \right)^{-1} = \frac{1}{\Theta(G)} \prod_{\chi\in\widehat{G}\setminus\{\chi_1\}} \left(\sum_{g\in G} \chi(g)x_g g \right).$$

1.2. Main results for dihedral groups and generalized quaternion groups. We give an analog of Theorem 1 on the group algebra for dihedral groups and generalized quaternion groups. Furthermore, we describe the group determinant of these groups as a circulant determinant of homogeneous polynomials. Since the expression for the general case is troublesome, we illustrate the result on a dihedral group $D_3 = \{e, a, a^2, b, ab, a^2b\}$. Let $\langle a\rangle$ be the subgroup generated by $\{a\}$ and A_h be a homogeneous polynomial defined by

$$A_h = \sum_{g\in\langle a\rangle} \chi_2(g)x_g x_{hg}$$

for $h \in G$. We describe $\Theta(D_3)$ as a circulant determinant of A_h.

Theorem 4. *Let G be D_3. Then, we have*

$$\Theta(D_3) = \prod_{\chi\in\widehat{\langle a\rangle}} \sum_{g\in\langle a\rangle} \chi(g)A_g.$$

Furthermore, we obtain the analog of Theorem 1 on the group algebra for D_3.

Theorem 5. *Let G be D_3, ω be a primitive third roots of unity and χ_2 be the non-trivial degree one representation. Then, we have*

$$\begin{aligned}\Theta(D_3)e &= \alpha_1\alpha_2 \left(A_e e + \omega A_a a + \omega^2 A_{a^2} a^2\right)\left(A_e e + \omega^2 A_a a + \omega A_{a^2} a^2\right)\\ &= \prod_{\chi\in\widehat{\langle a\rangle}} \sum_{g\in\langle a\rangle} \chi(g)A_g g,\end{aligned}$$

where

$$\alpha_1 = \sum_{g\in D_3} x_g g, \quad \alpha_2 = \sum_{g\in D_3} \chi_2(g)x_{g^{-1}} g.$$

Theorem 1 for D_3 and Theorem 4 follow by applying F to Theorem 5. Moreover, we obtain the following formula for inverse elements in the group algebra for D_3.

Corollary 6. *Let G be D_3. If $\Theta(G) \neq 0$, then we have*

$$\alpha_1^{-1} = \frac{1}{\Theta(D_3)}\alpha_2 \left(A_e e + \omega A_a a + \omega^2 A_{a^2} a^2\right) \left(A_e e + \omega^2 A_a a + \omega A_{a^2} a^2\right).$$

We can extend the results in Section 1.2 into dihedral groups and generalized quaternion groups.

1.3. Algebraic properties of irreducible factors corresponding to degree one representations. Let G be a dihedral group D_m or generalized quaternion group Q_m. The commutators of irreducible factors of the factorization of the group determinant on the group algebra corresponding to degree one representations have the following algebraic properties.

Theorem 7. *Let G be D_m or Q_m. Then, we have*

$$[\alpha_1, \alpha_2] = 0, \quad [\alpha_3, \alpha_4] = 0,$$

$$[\alpha_1, \alpha_3 + \alpha_4] = 0, \quad [\alpha_2, \alpha_3 + \alpha_4] = 0, \quad [\alpha_3, \alpha_1 + \alpha_2] = 0, \quad [\alpha_4, \alpha_1 + \alpha_2] = 0.$$

where $\alpha_i (1 \leq i \leq 4)$ is the irreducible factor of the factorization of the group determinant on the group algebra corresponding to the degree one representation. However, the equations for α_3 and α_4 do not hold when $G = D_m$ and m is odd.

We remark that when m is odd the number of degree one representation of D_m is two and when m is even the number is four. The number of degree one representation of Q_m is always four. From Theorem 7, we know that the degree one representations form natural pairing (χ_1, χ_2) and (χ_3, χ_4).

1.4. Future works. From the above results and Theorem 1, we expect to find determinant formulas for matrices with elements on the group algebra.

2. Factorization of group determinant

In this section, we recall the group determinant and the factorization of the group determinant.

2.1. Factorization of group determinant. Let G be a finite group of order n and $\{x_g | g \in G\}$ be independent commuting variables. The group determinant $\Theta(G)$ is the determinant of the $n \times n$ matrix $(x_{g,h})$ where $x_{g,h} = x_{gh^{-1}}$ for $g, h \in G$ and is thus a homogeneous polynomial of degree n in x_g. Frobenius gave the following theorem about the factorization of the group determinant.

Theorem 8 (Frobenius)**.** *Let $\widehat{G}$ be a complete set of irreducible representations of G. Then, we have*

$$\Theta(G) = \prod_{\varphi \in \widehat{G}} \det \left(\sum_{g \in G} \varphi(g) x_g \right)^{\deg \varphi}.$$

3. Results for finite abelian groups

In this section, we give an analog of Frobenius' theorem for finite abelian groups about factorization of the group determinant on the group algebra.

3.1. Preparation for main results of finite abelian groups. We prepare to explain for main results of finite abelian groups. Let $\{\varphi^{(1)}, \varphi^{(2)}, \ldots, \varphi^{(s)}\}$ be a complete set of irreducible representations of G, $M_k(\mathbb{C})$ be a set of $k \times k$ matrices with entries in $\mathbb{C}$ and $L(G)$ be the linear space of all complex functions on G. The set $L(G)$ is a ring with addition taken pointwise and convolution as multiplication [1, Theorem 5.2.3].

Definition 9 (Fourier transform)**.** *Define*

$$T : L(G) \to M_{d_1}(\mathbb{C}) \times M_{d_2}(\mathbb{C}) \times \cdots \times M_{d_s}(\mathbb{C})$$

by

$$Tf = (\widehat{f}(\varphi^{(1)}), \widehat{f}(\varphi^{(2)}), \ldots, \widehat{f}(\varphi^{(s)})).$$

where

$$\widehat{f}(\varphi^{(k)}) = \sum_{g \in G} \overline{\varphi^{(k)}(g)} f(g),$$

the bar denotes complex conjugation. We call Tf the Fourier transform of f.

Theorem 10 ([1], Theorem 5.5.6)**.** *The Fourier transform*

$$T : L(G) \to M_{d_1}(\mathbb{C}) \times M_{d_2}(\mathbb{C}) \times \cdots \times M_{d_s}(\mathbb{C})$$

is an isomorphism of rings.

Let

$$\mathbb{C}G = \left\{ \sum_{g \in G} x_g g \mid x_g \in \mathbb{C}, g \in G \right\}$$

be the group algebra of G.

Lemma 11. *Let G be a finite group. Then, we have*

$$L(G) \cong \mathbb{C}G.$$

as an isomorphism of rigns.

3.2. Results for finite abelian groups. We give the following theorem that is an analog of Theorem 8 for finite abelian groups on the group algebra.

Theorem 12. *Let G be a finite abelian group and e be the unit element of G. Then, we have*

$$\Theta(G)e = \prod_{\chi \in \widehat{G}} \sum_{g \in G} \chi(g) x_g g.$$

Proof. Let $G = \{g_1, g_2, \ldots, g_n\}$ and $\widehat{G} = \{\chi_1, \chi_2, \ldots, \chi_n\}$. By definition, we have

$$\alpha = \sum_{i=1}^{n} x_{g_i} g_i \in \mathbb{C}G.$$

Then, for $\chi_k \in \widehat{G}$, we define

$$\alpha_{\chi_k} = \sum_{i=1}^{n} \chi_k(g_i) x_{g_i} g_i \in \mathbb{C}G, \quad \alpha^*_{\chi_k} = \sum_{i=1}^{n} \chi_k(g_i) x_{g_i} \in \mathbb{C}.$$

We can regard α_{χ_k} as an element in $L(G)$ by Lemma 11, then we have

$$\alpha_{\chi_k}(g_i) = \chi_k(g_i) x_{g_i}.$$

The l-th component of the Fourier transform T of α_{χ_k} is

$$\begin{aligned}\widehat{\alpha_{\chi_k}}(\chi_l) &= \sum_{i=1}^{n} \overline{\chi_l(g_i)}\alpha_{\chi_k}(g_i) \\ &= \sum_{i=1}^{n} \overline{\chi_l(g_i)}\chi_k(g_i)x_{g_i} \\ &= \sum_{i=1}^{n} (\chi_k(\chi_l)^{-1})(g_i)x_{g_i} \\ &= \alpha^*_{\chi_k\chi_l^{-1}}.\end{aligned}$$

Therefore, by Theorem 8, we have

$$\begin{aligned}T(\alpha_{\chi_1})T(\alpha_{\chi_2})\cdots T(\alpha_{\chi_n}) &= \alpha^*_{\chi_1}\alpha^*_{\chi_2}\cdots\alpha^*_{\chi_n}(1,1,\ldots,1) \\ &= \Theta(G)(1,1,\ldots,1).\end{aligned}$$

By Theorem 10, we have

$$\alpha_{\chi_1}\alpha_{\chi_2}\cdots\alpha_{\chi_n} = \Theta(G)e.$$

□

Theorem 12 is stronger than Theorem 8 for finite abelian groups. In fact, let $F : \mathbb{C}G \to \mathbb{C}$ be a $\mathbb{C}$-linear ring homomorphism such that $F(g) = 1$ for all $g \in G$, Theorem 8 for finite abelian groups follows by applying F to Theorem 12. Moreover, we obtain the following formula for inverse elements in the group algebra.

Corollary 13. *Let G be a finite abelian group and χ_1 be the trivial representation of G. If $\Theta(G) \neq 0$, then we have*

$$\left(\sum_{g\in G} x_g g\right)^{-1} = \frac{1}{\Theta(G)} \prod_{\chi\in\widehat{G}\setminus\{\chi_1\}} \left(\sum_{g\in G} \chi(g)x_g g\right).$$

4. Recall dihedral groups and generalized quaternion groups

In this section, we recall dihedral groups and generalized quaternion groups.

4.1. Recall dihedral groups. We recall dihedral group D_m given by the presentation

$$D_m = \langle a, b \mid a^m = e, b^2 = e, b^{-1}ab = a^{-1}\rangle.$$

Lemma 14. *Every element $g \in D_m$ can be written uniquely as $g = a^k b^l$ where $0 \leq k < m$, and $l = 0, 1$. Namely, the order of D_m is $2m$.*

We have the following list of irreducible representations of D_m([2, Theorem 3]). Let ω be a primitive m-th roots of unity, for $1 \leq k \leq m-1$.

(1) When m is odd, and $1 \leq l \leq \frac{m-1}{2}$.

	e	a^k	b	a^kb
χ_1	1	1	1	1
χ_2	1	1	-1	-1
φ_l	$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$	$\begin{bmatrix} \omega^{lk} & 0 \\ 0 & \omega^{-lk} \end{bmatrix}$	$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \omega^{lk} \\ \omega^{-lk} & 0 \end{bmatrix}$

(2) When m is even, and $1 \leq l \leq \frac{m}{2} - 1$.

	e	a^k	b	a^kb
χ_1	1	1	1	1
χ_2	1	1	-1	-1
χ_3	1	$(-1)^k$	1	$(-1)^k$
χ_4	1	$(-1)^k$	-1	$(-1)^{k+1}$
φ_l	$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$	$\begin{bmatrix} \omega^{lk} & 0 \\ 0 & \omega^{-lk} \end{bmatrix}$	$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \omega^{lk} \\ \omega^{-lk} & 0 \end{bmatrix}$

4.2. Recall generalized quaternion groups. We recall generalized quaternion group Q_m given by the presentation

$$Q_m = \langle a, b \mid a^{2m} = e, b^2 = a^m, b^{-1}ab = a^{-1} \rangle.$$

Lemma 15. *Every element $g \in Q_m$ can be written uniquely as $g = a^k b^l$ where $0 \leq k < 2m$, and $l = 0, 1$. Namely, the order of Q_m is $4m$.*

We have the following list of irreducible representations of Q_m. Let ω be a primitive $2m$-th roots of unity, for $1 \leq k \leq 2m-1$.

(1) When m is odd, and $1 \leq l \leq m-1$.

	e	a^k	b	a^kb
χ_1	1	1	1	1
χ_2	1	1	-1	-1
χ_3	1	$(-1)^k$	i	$i(-1)^k$
χ_4	1	$(-1)^k$	$-i$	$i(-1)^{k+1}$
φ_l	$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$	$\begin{bmatrix} \omega^{lk} & 0 \\ 0 & \omega^{-lk} \end{bmatrix}$	$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \omega^{lk} \\ -\omega^{-lk} & 0 \end{bmatrix}$

(2) When m is even, and $1 \leq l \leq m-1$.

	e	a^k	b	a^kb
χ_1	1	1	1	1
χ_2	1	1	-1	-1
χ_3	1	$(-1)^k$	1	$(-1)^k$
χ_4	1	$(-1)^k$	-1	$(-1)^{k+1}$
φ_l	$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$	$\begin{bmatrix} \omega^{lk} & 0 \\ 0 & \omega^{-lk} \end{bmatrix}$	$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \omega^{lk} \\ -\omega^{-lk} & 0 \end{bmatrix}$

5. Main results for dihedral groups and generalized quaternion groups

In this section, we describe the group determinant of dihedral groups and generalized quaternion groups as a circulant determinant of homogeneous polynomials and we give an analog of Frobenius' theorem for dihedral groups and generalized quaternion groups about factorization of the group determinant on the group algebra.

5.1. Describe some group determinant as a circulant determinant. Let G be D_m or Q_m, $\langle a\rangle$ be the subgroup generated by $\{a\}$ and define

$$\chi'_l(a^k) = \omega^{lk}$$

for $0 \leq l \leq |\langle a\rangle| - 1$ where $|H|$ is the order of the group H. Notice that χ'_l is a degree one representation of $\langle a\rangle$. Let A_h be a homogeneous polynomial defined by

$$A_h = \sum_{g\in\langle a\rangle} \left(x_g x_{hg} - x_{gb} x_{hgb^{-1}}\right)$$

for $h \in G$. This homogeneous polynomial A_h has the following properties.

Lemma 16. *Let G be D_m or Q_m. Then, we have*

(1) *If $h \in G \setminus \langle a\rangle$, $A_h = 0$.*
(2) *For all $h \in G$, $A_h = A_{h^{-1}}$.*

Proof. Element $h \in G \setminus \langle a\rangle$ can be written as $a^k b$ for $0 \leq k \leq |\langle a\rangle| - 1$. If $g \in \langle a\rangle$, then $bg = g^{-1}b$ and we can write g as $a^k g^{-1}$. From these, we have

$$\begin{aligned}
A_{a^k b} &= \sum_{g\in\langle a\rangle} x_g x_{a^k bg} - \sum_{g\in\langle a\rangle} x_{gb} x_{a^k bgb^{-1}} \\
&= \sum_{g\in\langle a\rangle} x_{a^k g^{-1}} x_{a^k b a^k g^{-1}} - \sum_{g\in\langle a\rangle} x_{gb} x_{a^k g^{-1}} \\
&= \sum_{g\in\langle a\rangle} x_{a^k g^{-1}} x_{gb} - \sum_{g\in\langle a\rangle} x_{a^k g^{-1}} x_{gb} \\
&= 0.
\end{aligned}$$

This proves the first claim. If $h \in G \setminus \langle a\rangle$, then, we have $h^{-1} \in G \setminus \langle a\rangle$. (2) follows from first claim. We assume that $h \in \langle a\rangle$. We can write g as $h^{-1}g$ in the first sum and g as $g = h^{-1}gb^{-2}$ in the second. From this and $b^{-3} = b$, we have

$$\begin{aligned}
A_h &= \sum_{g\in\langle a\rangle} x_g x_{hg} - \sum_{g\in\langle a\rangle} x_{gb} x_{hgb^{-1}} \\
&= \sum_{g\in\langle a\rangle} x_{(h^{-1}g)} x_{h(h^{-1}g)} - \sum_{g\in\langle a\rangle} x_{(h^{-1}gb^{-2})b} x_{h(h^{-1}gb^{-2})b^{-1}} \\
&= \sum_{g\in\langle a\rangle} x_{h^{-1}g} x_g - \sum_{g\in\langle a\rangle} x_{h^{-1}gb^{-1}} x_{gb^{-3}} \\
&= \sum_{g\in\langle a\rangle} x_g x_{h^{-1}g} - \sum_{g\in\langle a\rangle} x_{h^{-1}gb^{-1}} x_{gb} \\
&= A_{h^{-1}}.
\end{aligned}$$

This completes the proof. □

Lemma 17. *Let G be D_m or Q_m and $\chi_l (1 \leq l \leq 4)$ be a degree one representation of Section 4. Then, we have*

$$\sum_{g\in G} \chi_1(g)x_g \sum_{g'\in G} \chi_2(g')x_{g'} = \sum_{h\in\langle a\rangle} A_h,$$
$$\sum_{g\in G} \chi_3(g)x_g \sum_{g'\in G} \chi_4(g')x_{g'} = \sum_{h\in\langle a\rangle} \chi'_{\frac{|\langle a\rangle|}{2}}(h)A_h.$$

This lemma will be proved later.

Lemma 18. *Let G be D_m or Q_m and φ_l be degree two representations of Section 4. Then, we have*

$$\det\left(\sum_{g\in G} \varphi_l(g)x_g\right) = \sum_{h\in\langle a\rangle} \chi'_l(h)A_h.$$

Proof. Define the function δ by

$$\delta(G) = \begin{cases} 1 & G = D_m, \\ -1 & G = Q_m. \end{cases}$$

We obtain

$$\begin{aligned}
\det\left(\sum_{g\in G} \varphi_l(g)x_g\right) &= \det\left(\sum_{g\in\langle a\rangle} \varphi_l(g)x_g + \sum_{g\in G\setminus\langle a\rangle} \varphi_l(g)x_g\right) \\
&= \det\left(\sum_{g\in\langle a\rangle} \varphi_l(g)x_g + \sum_{g\in\langle a\rangle} \varphi_l(gb)x_{gb}\right) \\
&= \det\begin{bmatrix} \sum_{g\in\langle a\rangle} \chi'_l(g)x_g & \sum_{g\in\langle a\rangle} \chi'_l(g)x_{gb} \\ \delta(G)\sum_{g\in\langle a\rangle} \chi'_l(g^{-1})x_{gb} & \sum_{g\in\langle a\rangle} \chi'_l(g^{-1})x_g \end{bmatrix} \\
&= \sum_{h\in\langle a\rangle}\sum_{g\in\langle a\rangle} \chi'_l(hg^{-1})x_h x_g - \delta(G)\sum_{g\in\langle a\rangle}\sum_{h\in\langle a\rangle} \chi'_l(g^{-1}h)x_{gb}x_{hb}. \qquad (1)
\end{aligned}$$

We can write h as hg in the first sum and h as hgb^{-2} in the second. Then (1) equals

$$\begin{aligned}
&\sum_{h\in\langle a\rangle} \chi'_l(h)\sum_{g\in\langle a\rangle} x_g x_{hg} - \delta(G)\sum_{g\in\langle a\rangle}\sum_{h\in\langle a\rangle} \chi'_l(g^{-1}(hgb^{-2}))x_{gb}x_{(hgb^{-2})b} \\
&= \sum_{h\in\langle a\rangle} \chi'_l(h)\sum_{g\in\langle a\rangle} x_g x_{hg} - \delta(G)\chi'_l(b^{-2})\sum_{h\in\langle a\rangle} \chi'_l(h)\sum_{g\in\langle a\rangle} x_{gb}x_{hgb^{-1}}. \qquad (2)
\end{aligned}$$

It is easy to see that $\delta(G)\chi'_l(b^{-2}) = 1$. Then, (2) equals

$$\sum_{h\in\langle a\rangle} \chi'_l(h)\left(\sum_{g\in\langle a\rangle} (x_g x_{hg} - x_{gb}x_{hgb^{-1}})\right) = \sum_{h\in\langle a\rangle} \chi'_l(h)A_h$$

as required. □

Lemma 19. *Let χ' be a degree one representation of $\langle a\rangle$. Then, we have*

$$\sum_{g\in\langle a\rangle}\chi'(g)A_g=\sum_{g\in\langle a\rangle}\overline{\chi'(g)}A_g.$$

Proof. This follows from Lemma 16. □

Let $\widehat{H}$ be a complete set of irreducible representations of the group H. We describe $\Theta(G)$ as a circulant determinant of A_h.

Theorem 20. *Let G be D_m or Q_m. Then, we have*

$$\Theta(G)=\prod_{\chi'\in\widehat{\langle a\rangle}}\sum_{g\in\langle a\rangle}\chi'(g)A_g.$$

Proof. First, let $G=D_m$ and m be odd. By Theorem 8 and Lemma 17, we have

$$\Theta(G)=\left(\sum_{g\in\langle a\rangle}A_g\right)\prod_{l=1}^{\frac{m-1}{2}}\left(\sum_{g\in\langle a\rangle}\chi'_l(g)A_g\right)^2.$$

From Lemma 19, we have

$$\begin{aligned}\Theta(G)&=\left(\sum_{g\in\langle a\rangle}A_g\right)\prod_{l=1}^{\frac{m-1}{2}}\left(\sum_{g\in\langle a\rangle}\chi'_l(g)A_g\right)\left(\sum_{g\in\langle a\rangle}\chi'_{m-l}(g)A_g\right)\\&=\prod_{l=0}^{m-1}\sum_{g\in\langle a\rangle}\chi'_l(g)A_g.\end{aligned}$$

In other case, by Theorem 8 and Lemma 17, we have

$$\Theta(G)=\left(\sum_{g\in\langle a\rangle}A_g\right)\left(\sum_{g\in\langle a\rangle}\chi'_{\frac{|\langle a\rangle|}{2}}(g)A_g\right)\prod_{l=1}^{\frac{|\langle a\rangle|}{2}-1}\left(\sum_{g\in\langle a\rangle}\chi'_l(g)A_g\right)^2.$$

From Lemma 19, we have

$$\begin{aligned}\Theta(G)&=\left(\sum_{g\in\langle a\rangle}A_g\right)\left(\sum_{g\in\langle a\rangle}\chi'_{\frac{|\langle a\rangle|}{2}}(g)A_g\right)\\&\quad\prod_{l=1}^{\frac{|\langle a\rangle|}{2}-1}\left(\sum_{g\in\langle a\rangle}\chi'_l(g)A_g\right)\left(\sum_{g\in\langle a\rangle}\chi'_{|\langle a\rangle|-l}(g)A_g\right)\\&=\prod_{l=0}^{|\langle a\rangle|-1}\sum_{g\in\langle a\rangle}\chi'_l(g)A_g.\end{aligned}$$

□

5.2. Main results for dihedral groups and generalized quaternion groups. Let G be D_m or Q_m and $\chi_l (1 \leq l \leq 4)$ be a degree one representation of Section 4. By definition, we have

$$\alpha_1 = \sum_{g \in G} x_g g, \quad \alpha_2 = \sum_{g \in \langle a \rangle} \chi_2(g) x_{g^{-1}} g + \sum_{g \in G \setminus \langle a \rangle} \chi_2(g) x_g g,$$
$$\alpha_3 = \sum_{g \in G} \chi_3(g) x_g g, \quad \alpha_4 = \sum_{g \in \langle a \rangle} \chi_4(g) x_{g^{-1}} g + \sum_{g \in G \setminus \langle a \rangle} \chi_4(g) x_g g.$$

Lemma 21. *The following formula holds.*

$$\alpha_1 \alpha_2 = \sum_{h \in \langle a \rangle} A_h h.$$

Proof. We have

$$\begin{aligned}
\alpha_1 \alpha_2 &= \sum_{g' \in G} x_{g'} g' \sum_{g \in \langle a \rangle} \chi_2(g) x_{g^{-1}} x_{g^{-1}} g + \sum_{g' \in G} x_{g'} g' \sum_{g \in G \setminus \langle a \rangle} \chi_2(g) x_g g \\
&= \sum_{g' \in G} \sum_{g \in \langle a \rangle} \chi_2(g) x_{g'} x_{g^{-1}} g' g + \sum_{g' \in G} \sum_{g \in G \setminus \langle a \rangle} \chi_2(g) x_{g'} x_g g' g \\
&= \sum_{g' \in G} \sum_{g \in \langle a \rangle} x_{g'} x_{g^{-1}} g' g + \sum_{g' \in G} \sum_{g \in \langle a \rangle} \chi_2(gb) x_{g'} x_{gb} g' g b \\
&= \sum_{g' \in G} \sum_{g \in \langle a \rangle} x_{g'} x_{g^{-1}} g' g - \sum_{g' \in G} \sum_{g \in \langle a \rangle} x_{g'} x_{gb} g' g b \\
&= \sum_{g' \in G} \sum_{g \in \langle a \rangle} x_{g'} x_{g^{-1}} g' g - \sum_{g' \in G} \sum_{g \in \langle a \rangle} x_{g'} x_{gb} g' b g^{-1} \\
&= \sum_{g' \in G} \sum_{g \in \langle a \rangle} x_{g'} x_{g^{-1}} g' g - \sum_{g' \in G} \sum_{g \in \langle a \rangle} x_{g' b^{-1}} x_{gb} g' g^{-1} \\
&= \sum_{g' \in G} \sum_{g \in \langle a \rangle} x_{g'} x_{g^{-1}} g' g - \sum_{g' \in G} \sum_{g \in \langle a \rangle} x_{g' b^{-1}} x_{g^{-1} b} g' g. \qquad (3)
\end{aligned}$$

We can make the change of variables $h = g'g$ to compute, then (3) equals

$$\begin{aligned}
&\sum_{h \in G} \sum_{g \in \langle a \rangle} x_{hg^{-1}} x_{g^{-1}} h - \sum_{h \in G} \sum_{g \in \langle a \rangle} x_{hg^{-1} b^{-1}} x_{g^{-1} b} h \\
&= \sum_{h \in G} \sum_{g \in \langle a \rangle} x_{hg} x_g h - \sum_{h \in G} \sum_{g \in \langle a \rangle} x_{hgb^{-1}} x_{gb} h \\
&= \sum_{h \in G} \sum_{g \in \langle a \rangle} (x_g x_{hg} - x_{gb} x_{hgb^{-1}}) h \\
&= \sum_{h \in G} A_h h \\
&= \sum_{h \in \langle a \rangle} A_h h
\end{aligned}$$

as required. □

Lemma 22. *The following formula holds.*

$$\alpha_3\alpha_4 = \sum_{h\in\langle a\rangle} \chi'_{\frac{|\langle a\rangle|}{2}}(h)A_h h.$$

Proof. We have

$$\begin{aligned}\alpha_3\alpha_4 &= \sum_{g'\in G}\chi_3(g')x_{g'}g'\sum_{g\in\langle a\rangle}\chi_4(g)x_{g^{-1}}g + \sum_{g'\in G}\chi_3(g')x_{g'}g'\sum_{g\in G\setminus\langle a\rangle}\chi_4(g)x_g g\\ &= \sum_{g'\in G}\sum_{g\in\langle a\rangle}\chi_3(g')\chi_4(g)x_{g'}x_{g^{-1}}g'g + \sum_{g'\in G}\sum_{g\in G\setminus\langle a\rangle}\chi_3(g')\chi_4(g)x_{g'}x_g g'g.\end{aligned}$$

If $g\in\langle a\rangle$, then, we have $\chi_3(g)\chi_4(g^{-1}) = 1$. We have

$$\begin{aligned}\sum_{g'\in G}\sum_{g\in\langle a\rangle}\chi_3(g')\chi_4(g)x_{g'}x_{g^{-1}}g'g &= \sum_{h\in G}\sum_{g\in\langle a\rangle}\chi_3(hg^{-1})\chi_4(g)x_{hg^{-1}}x_{g^{-1}}h\\ &= \sum_{h\in G}\sum_{g\in\langle a\rangle}\chi_3(hg)\chi_4(g^{-1})x_{hg}x_g h\\ &= \sum_{h\in G}\sum_{g\in\langle a\rangle}\chi_3(h)x_g x_{hg}h.\end{aligned}$$

We have

$$\begin{aligned}\sum_{g'\in G}\sum_{g\in G\setminus\langle a\rangle}\chi_3(g')\chi_4(g)x_{g'}x_g g'g &= \sum_{g'\in G}\sum_{g\in\langle a\rangle}\chi_3(g')\chi_4(gb)x_{g'}x_{gb}g'gb\\ &= \chi_4(b)\sum_{g'\in G}\sum_{g\in\langle a\rangle}\chi_3(g')\chi_4(g)x_{g'}x_{gb}g'bg^{-1}\\ &= \chi_4(b)\sum_{g'\in G}\sum_{g\in\langle a\rangle}\chi_3(g'b^{-1})\chi_4(g)x_{g'b^{-1}}x_{gb}g'g^{-1}. \qquad (4)\end{aligned}$$

It is easy that to see $\chi_3(b^{-1})\chi_4(b) = -1$, then (4) equals

$$\begin{aligned}&-\sum_{g'\in G}\sum_{g\in\langle a\rangle}\chi_3(g')\chi_4(g)x_{g'b^{-1}}x_{gb}g'g^{-1}\\ =&-\sum_{h\in G}\sum_{g\in\langle a\rangle}\chi_3(hg^{-1})\chi_4(g^{-1})x_{hg^{-1}b^{-1}}x_{g^{-1}b}h\\ =&-\sum_{h\in G}\sum_{g\in\langle a\rangle}\chi_3(hg)\chi_4(g)x_{hgb^{-1}}x_{gb}h. \qquad (5)\end{aligned}$$

If $g\in\langle a\rangle$, then we have $\chi_3(g)\chi_4(g) = 1$ and (5) equals

$$-\sum_{h\in G}\sum_{g\in\langle a\rangle}\chi_3(h)x_{hgb^{-1}}x_{gb}h.$$

We compute

$$\begin{aligned}\alpha_3\alpha_4 &= \sum_{h\in G}\chi_3(h)\sum_{g\in\langle a\rangle}(x_g x_{hg} - x_{gb}x_{hgb^{-1}})h\\ &= \sum_{h\in G}\chi_3(h)A_h h\\ &= \sum_{h\in\langle a\rangle}\chi_3(h)A_h h\\ &= \sum_{h\in\langle a\rangle}\chi'_{\frac{|\langle a\rangle|}{2}}(h)A_h h\end{aligned}$$

as required. □

Lemmas 21 and 22 are stronger than Lemma 17. In fact, Lemma 17 follows by applying F to Lemmas 21 and 22.

Theorem 23. *Let G be D_m or Q_m and e be the unit element of G. Then, we have*

$$\Theta(G)e = \prod_{\chi'\in\widehat{\langle a\rangle}}\sum_{g\in\langle a\rangle}\chi'(g)x_g g.$$

Proof. The group $\langle a\rangle$ is a finite abelian group and by Theorem 12, there exists a C such that

$$Ce = \prod_{\chi'\in\widehat{\langle a\rangle}}\sum_{g\in\langle a\rangle}\chi'(g)x_g g.$$

By Theorem 20 and the mapping F, we have

$$C = \Theta(G).$$

This completes the proof. □

Theorem 8 for D_m or Q_m and Theorem 20 follow by applying F to Theorem 23. Moreover, we obtain the following formula for inverse elements in the group algebra for D_m and Q_m.

Corollary 24. *Let G be D_m or Q_m. If $\Theta(G)\neq 0$,*

(1) *When $G = D_m$ and m is odd, we have*

$$\alpha_1^{-1} = \frac{1}{\Theta(G)}\alpha_2\prod_{\chi'\in\widehat{\langle a\rangle}\setminus\{\chi'_0\}}\sum_{g\in\langle a\rangle}\chi'(g)x_g g.$$

(2) *In other cases, we have*

$$\alpha_1^{-1} = \frac{1}{\Theta(G)}\alpha_2\alpha_3\alpha_4\prod_{\chi'\in\widehat{\langle a\rangle}\setminus\{\chi'_0\}}\sum_{g\in\langle a\rangle}\chi'(g)x_g g.$$

Note that the group algebra is non-commutative, so the order of the factors is important.

6. Algebraic properties of irreducible factors corresponding to degree one representations

In this section, we follow the commutators of irreducible factors of the factorization of the group determinant on the group algebra for D_m and Q_m corresponding to degree one representations have interesting algebraic properties. From this result, we know that degree one representations from natural pairing.

6.1. Transformations of irreducible factors corresponding to degree one representations. In this subsection, we obtain transformations of irreducible factors corresponding to degree one representations. We need it to prove for algebraic properties of irreducible factors corresponding to degree one representations.

Lemma 25. *Let G be D_m or Q_m. Under the change of variable*

$$x_g \mapsto \begin{cases} \chi_2(g)x_{g^{-1}} & \text{if} \quad g \in \langle a \rangle, \\ \chi_2(g)x_g & \text{otherwise.} \end{cases}$$

Then α_1 becomes α_2 by and vice versa, the same holds for α_3 and α_4.

Proof. From the definition of α_1, the factor α_1 becomes α_3 under the change of variable. The factor α_2 becomes

$$\begin{aligned} \sum_{g\in\langle a\rangle} \chi_2(g)\chi_2(g^{-1})x_g g + \sum_{g\in G\setminus\langle a\rangle} \chi_2(g)\chi_2(g)x_g g &= \sum_{g\in\langle a\rangle} x_g g + \chi_2(b)^2 \sum_{g\in G\setminus\langle a\rangle} x_g g \\ &= \sum_{g\in\langle a\rangle} x_g g + \sum_{g\in G\setminus\langle a\rangle} x_g g \\ &= \sum_{g\in G} x_g g \\ &= \alpha_1. \end{aligned}$$

The factor α_3 becomes

$$\begin{aligned} &\sum_{g\in\langle a\rangle} \chi_3(g)\chi_2(g)x_{g^{-1}}g + \sum_{g\in G\setminus\langle a\rangle} \chi_3(g)\chi_2(g)x_g g \\ &= \sum_{g\in\langle a\rangle} \chi_3(g)x_{g^{-1}}g - \sum_{g\in G\setminus\langle a\rangle} \chi_3(g)x_g g \\ &= \sum_{g\in\langle a\rangle} \chi_4(g)x_{g^{-1}}g + \sum_{g\in G\setminus\langle a\rangle} \chi_4(g)x_g g \\ &= \alpha_4. \end{aligned}$$

The factor α_4 becomes

$$\begin{aligned} &\sum_{g\in\langle a\rangle} \chi_4(g)\chi_2(g^-1)x_g g + \sum_{g\in G\setminus\langle a\rangle} \chi_4(g)\chi_2(g)x_g g \\ &= \sum_{g\in\langle a\rangle} \chi_4(g)x_g g - \sum_{g\in G\setminus\langle a\rangle} \chi_4(g)x_g g \\ &= \sum_{g\in\langle a\rangle} \chi_3(g)x_g g + \sum_{g\in G\setminus\langle a\rangle} \chi_3(g)x_g g \\ &= \alpha_3. \end{aligned}$$

This completes the proof. □

Lemma 26. *Let G be D_m or Q_m. Under the change of variable $x_g \mapsto \chi_3(g)x_g$, the factor α_1 becomes α_3, the factor α_2 becomes α_4, and α_3 becomes*

$$\sum_{g\in\langle a\rangle} x_g g - \sum_{g\in G\setminus\langle a\rangle} x_g g$$

in the case that m is odd and α_1 otherwise. The factor α_4 becomes

$$\sum_{g\in\langle a\rangle} x_{g^{-1}} g + \sum_{g\in G\setminus\langle a\rangle} x_g g$$

in the case that m is odd and α_2 otherwise. Namely, the factor $\alpha_3+\alpha_4$ becomes $\alpha_1+\alpha_2$.

Proof. In the definition of α_1, we obtain α_3. The factor α_2 becomes

$$\begin{aligned}
&\sum_{g\in\langle a\rangle} \chi_2(g)\chi_3(g^{-1})x_{g^{-1}}g + \sum_{g\in G\setminus\langle a\rangle} \chi_2(g)\chi_3(g)x_g g \\
&\quad= \sum_{g\in\langle a\rangle} \chi_3(g)x_{g^{-1}}g - \sum_{g\in G\setminus\langle a\rangle} \chi_3(g)x_g g \\
&\quad= \sum_{g\in\langle a\rangle} \chi_4(g)x_{g^{-1}}g + \sum_{g\in G\setminus\langle a\rangle} \chi_4(g)x_g g \\
&\quad= \alpha_4.
\end{aligned}$$

The factor α_3 becomes

$$\sum_{g\in G} \chi_3(g)^2 x_g g = \sum_{g\in\langle a\rangle} x_g g + \chi_3(b)^2 \sum_{g\in G\setminus\langle a\rangle} x_g g.$$

When m is even, right hand side is equal to α_1, otherwise is equal to

$$\sum_{g\in\langle a\rangle} x_g g - \sum_{g\in G\setminus\langle a\rangle} x_g g.$$

The factor α_4 becomes

$$\begin{aligned}
&\sum_{g\in\langle a\rangle} \chi_4(g)\chi_3(g^{-1})x_{g^{-1}}g + \sum_{g\in G\setminus\langle a\rangle} \chi_4(g)\chi_3(g)x_g g \\
&\quad= \sum_{g\in\langle a\rangle} x_{g^{-1}}g - \chi_3(b)^2 \sum_{g\in G\setminus\langle a\rangle} x_g g,
\end{aligned}$$

When m is odd, right hand side is equal to α_2, otherwise is equal to

$$\sum_{g\in\langle a\rangle} x_{g^{-1}} g + \sum_{g\in G\setminus\langle a\rangle} x_g g.$$

This completes the proof. □

6.2. Algebraic properties of irreducible factors corresponding to degree one representations. In this subsection, we follow the commutators of irreducible factors of the factorization of the group determinant on the group algebra for D_m and Q_m corresponding to degree one representations have interesting algebraic properties. From this result, we know that degree one representations form natural pairing.

Lemma 27. *If $h \in \langle a \rangle$, then we have*

$$\sum_{g \in \langle a \rangle} x_{gb} x_{b^{-1}g^{-1}h} = \sum_{g \in \langle a \rangle} x_{gb} x_{hgb^{-1}}.$$

Proof. We compute

$$\begin{aligned}
\sum_{g \in \langle a \rangle} x_{gb} x_{b^{-1}g^{-1}h} &= \sum_{g \in \langle a \rangle} x_{b^{-1}g^{-1}h} x_{gb} \\
&= \sum_{g \in \langle a \rangle} x_{h^{-1}gb^{-1}} x_{gb} \\
&= \sum_{g \in \langle a \rangle} x_{h^{-1}gb^{3}} x_{gb} \\
&= \sum_{g \in \langle a \rangle} x_{h^{-1}(hgb^{-2})b^{3}} x_{hgb^{-2}b} \\
&= \sum_{g \in \langle a \rangle} x_{gb} x_{hgb^{-1}}
\end{aligned}$$

as required. □

Lemma 28. *If $h \in G \setminus \langle a \rangle$, then we have*

$$\sum_{g \in G \setminus \langle a \rangle} x_g x_{g^{-1}h} = \sum_{g \in \langle a \rangle} x_g x_{gh}.$$

Proof. We compute

$$\begin{aligned}
\sum_{g \in G \setminus \langle a \rangle} x_g x_{g^{-1}h} &= \sum_{g \in \langle a \rangle} x_{gb} x_{b^{-1}g^{-1}a^k b} \\
&= \sum_{g \in \langle a \rangle} x_{gb} x_{b^{-1}bga^{-k}} \\
&= \sum_{g \in \langle a \rangle} x_{gb} x_{ga^{-k}} \\
&= \sum_{g \in \langle a \rangle} x_{ga^k b} x_g \\
&= \sum_{g \in \langle a \rangle} x_g x_{ga^k b} \\
&= \sum_{g \in \langle a \rangle} x_g x_{gh}
\end{aligned}$$

as required. □

Define $[a, b] = ab - ba$ where $a, b \in \mathbb{C}G$. This is equal to zero if and only if a and b commute.

Theorem 29. *The following formula holds.*

$$[\alpha_1, \alpha_2] = 0.$$

Proof. We have

$$\begin{aligned}\alpha_2\alpha_1 &= \left(\sum_{g\in\langle a\rangle} x_{g^{-1}}g - \sum_{g\in G\setminus\langle a\rangle} x_g g\right)\left(\sum_{g'\in G} x_{g'}g'\right)\\ &= \sum_{g'\in G}\left(\sum_{g\in\langle a\rangle} x_{g^{-1}}x_{g'} - \sum_{g\in G\setminus\langle a\rangle} x_g x_{g'}gg'\right)\\ &= \sum_{h\in G}\left(\sum_{g\in\langle a\rangle} x_{g^{-1}}x_{g^{-1}h} - \sum_{g\in G\setminus\langle a\rangle} x_g x_{g^{-1}h}\right)h.\end{aligned}$$

By Lemmas 25 and 28, we have

$$\begin{aligned}\alpha_2\alpha_1 &= \sum_{h\in\langle a\rangle}\left(\sum_{g\in\langle a\rangle} x_g x_{hg} - \sum_{g\in\langle a\rangle} x_{gb}x_{hgb^{-1}}\right)\\ &= \alpha_1\alpha_2.\end{aligned}$$

This completes the proof. □

Theorem 30. *The following formula holds.*

$$[\alpha_3,\alpha_4] = 0.$$

Proof. This theorem follows from Lemma 26 and Theorem 29. □

Lemma 31. *The following formula holds.*

$$[\alpha_1,\alpha_3] = \sum_{h\in G}\sum_{g\in G}\chi_3(g^{-1}h)x_g x_{g^{-1}h}h - \sum_{h\in G}\sum_{g\in G}\chi_3(g)x_g x_{g^{-1}h}h.$$

Proof. We compute

$$\begin{aligned}[\alpha_1,\alpha_3] &= \sum_{g\in G} x_g g\sum_{g'\in G}\chi_3(g')x_{g'}g' - \sum_{g\in G}\chi_3(g)x_g g\sum_{g'\in G} x_{g'}g'\\ &= \sum_{g\in G}\sum_{g'\in G}\chi_3(g')x_g x_{g'}gg' - \sum_{g\in G}\sum_{g'\in G}\chi_3(g)x_g x_{g'}gg'\\ &= \sum_{g\in G}\sum_{h\in G}\chi_3(g^{-1}h)x_g x_{g^{-1}h}h - \sum_{g\in G}\sum_{h\in G}\chi_3(g)x_g x_{g^{-1}h}h\end{aligned}$$

as required. □

Lemma 32. *Let h be a^k with k odd. Then, we have*

$$\sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{g^{-1}h} = 0.$$

Proof. We compute

$$\begin{aligned}\sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{g^{-1}h} &= \sum_{g\in\langle a\rangle}\chi_3(g^{-1}h)x_{g^{-1}h}x_g\\ &= -\sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{g^{-1}h}\end{aligned}$$

as required. □

Lemma 33. *The coefficient of $h = a^k$ in $[\alpha_1, \alpha_3]$ is*

$$(-1)^k \sum_{g \in G \setminus \langle a \rangle} \chi_3(g^{-1}) x_g x_{g^{-1}h} - \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h}.$$

Proof. We have

$$\begin{aligned}
&\sum_{g \in G} \chi_3(g^{-1}h) x_g x_{g^{-1}h} - \sum_{g \in G} \chi_3(g) x_g x_{g^{-1}h} \\
&\quad = (-1)^k \sum_{g \in G} \chi_3(g^{-1}) x_g x_{g^{-1}h} - \sum_{g \in G} \chi_3(g) x_g x_{g^{-1}h} \\
&\quad = (-1)^k \sum_{g \in \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h} + (-1)^k \sum_{g \in G \setminus \langle a \rangle} \chi_3(g^{-1}) x_g x_{g^{-1}h} \\
&\qquad - \sum_{g \in \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h} - \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h}.
\end{aligned}$$

From Lemma 32, this completes the proof. □

By Lemma 33, we know that

$$[\alpha_1, \alpha_3] \neq 0.$$

From Lemma 25, we have

$$[\alpha_2, \alpha_4] \neq 0.$$

Namely, we have to be careful with the factors in Corollary 24.

Lemma 34. *The coefficient of $h = a^k b$ in $[\alpha_1, \alpha_3]$ is*

(1) *For the case that m is odd.*

$$\begin{cases} (1+i) \sum_{g \in \langle a \rangle} \chi_3(g) x_g x_{gh} - (1+i) \sum_{g \in \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h}, & \text{if } k \text{ is odd.} \\ (1-i) \sum_{g \in \langle a \rangle} \chi_3(g) x_g x_{gh} - (1-i) \sum_{g \in \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h}, & \text{if } k \text{ is even.} \end{cases}$$

(2) *For the case that m is even.*

$$\begin{cases} -2 \sum_{g \in G} \chi_3(g) x_g x_{g^{-1}h}, & \text{if } k \text{ is odd.} \\ 0, & \text{if } k \text{ is even.} \end{cases}$$

Proof. (1) For the case that m is odd, we compute

$$\begin{aligned}
&\sum_{g\in G}\chi_3(g^{-1}h)x_gx_{g^{-1}h}-\sum_{g\in G}\chi_3(g)x_gx_{g^{-1}h}\\
&=(-1)^ki\sum_{g\in G}\chi_3(g^{-1})x_gx_{g^{-1}h}-\sum_{g\in G}\chi_3(g)x_gx_{g^{-1}h}\\
&=(-1)^ki\sum_{g\in\langle a\rangle}\chi_3(g^{-1})x_gx_{g^{-1}h}+(-1)^ki\sum_{g\in G\setminus\langle a\rangle}\chi_3(g^{-1})x_gx_{g^{-1}h}\\
&\quad-\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}-\sum_{g\in G\setminus\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}\\
&=(-1)^ki\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}+(-1)^ki\sum_{g\in\langle a\rangle}\chi_3(b^{-1}g^{-1})x_{gb}x_{b^{-1}g^{-1}h}\\
&\quad-\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}-\sum_{g\in\langle a\rangle}\chi_3(gb)x_{gb}x_{b^{-1}g^{-1}h}\\
&=(-1)^ki\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}+(-1)^k\sum_{g\in\langle a\rangle}\chi_3(g)x_{gb}x_{a^{-k}g}\\
&\quad-\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}-i\sum_{g\in\langle a\rangle}\chi_3(g)x_{gb}x_{a^{-k}g}\\
&=(-1)^ki\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}+(-1)^k\sum_{g\in\langle a\rangle}\chi_3(ga^k)x_{ga^kg}x_g\\
&\quad-\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}-i\sum_{g\in\langle a\rangle}\chi_3(ga^k)x_{ga^kb}x_g\\
&=(-1)^ki\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}+\sum_{g\in\langle a\rangle}\chi_3(g)x_{gh}x_g\\
&\quad-\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}+(-1)^{k+1}i\sum_{g\in\langle a\rangle}\chi_3(g)x_{gh}x_g
\end{aligned}$$

as required.

(2) For the case that m is even, we compute

$$\begin{aligned}
&\sum_{g\in G}\chi_3(g^{-1}h)x_gx_{g^{-1}h}-\sum_{g\in G}\chi_3(g)x_gx_{g^{-1}h}\\
&\quad=(-1)^k\sum_{g\in G}\chi_3(g)x_gx_{g^{-1}h}-\sum_{g\in G}\chi_3(g)x_gx_{g^{-1}h}
\end{aligned}$$

as required.

□

Lemma 35. *The following formula holds.*

$$\begin{aligned}
[\alpha_1,\alpha_4]=&\sum_{h\in G}\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{hg}h-\sum_{h\in G}\sum_{g\in G\setminus\langle a\rangle}\chi_3(g)x_gx_{hg^{-1}}h\\
&-\sum_{h\in G}\sum_{g\in\langle a\rangle}\chi_3(g)x_gx_{gh}h+\sum_{h\in G}\sum_{g\in G\setminus\langle a\rangle}\chi_3(g)x_gx_{g^{-1}h}h.
\end{aligned}$$

Proof. We compute

$$\begin{aligned}
[\alpha_1, \alpha_4] &= \sum_{g' \in G} x_{g'} g' \left(\sum_{g \in \langle a \rangle} \chi_4(g) x_{g^{-1}} g + \sum_{g \in G \setminus \langle a \rangle} \chi_4(g) x_g g \right) \\
&\quad - \left(\sum_{g \in \langle a \rangle} \chi_4(g) x_{g^{-1}} g + \sum_{g \in G \setminus \langle a \rangle} \chi_4(g) x_g g \right) \sum_{g' \in G} x_{g'} g' \\
&= \sum_{g' \in G} \sum_{g \in \langle a \rangle} \chi_4(g) x_{g'} x_{g^{-1}} g' g + \sum_{g' \in G} \sum_{g \in G \setminus \langle a \rangle} \chi_4(g) x_{g'} x_g g' g \\
&\quad - \sum_{g \in \langle a \rangle} \sum_{g' \in G} \chi_4(g) x_{g^{-1}} x_{g'} g g' - \sum_{g \in G \setminus \langle a \rangle} \sum_{g' \in G} \chi_4(g) x_g x_{g'} g g' \\
&= \sum_{h \in G} \sum_{g \in \langle a \rangle} \chi_4(g) x_{hg^{-1}} x_{g^{-1}} h + \sum_{h \in G} \sum_{g \in G \setminus \langle a \rangle} \chi_4(g) x_{hg^{-1}} x_g h \\
&\quad - \sum_{g \in \langle a \rangle} \sum_{h \in G} \chi_4(g) x_{g^{-1}} x_{g^{-1}h} h - \sum_{g \in G \setminus \langle a \rangle} \sum_{h \in G} \chi_4(g) x_g x_{g^{-1}h} h \\
&= \sum_{h \in G} \sum_{g \in \langle a \rangle} \chi_3(g) x_{hg^{-1}} x_{g^{-1}} h - \sum_{h \in G} \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_{hg^{-1}} x_g h \\
&\quad - \sum_{g \in \langle a \rangle} \sum_{h \in G} \chi_3(g) x_{g^{-1}} x_{g^{-1}h} h + \sum_{g \in G \setminus \langle a \rangle} \sum_{h \in G} \chi_3(g) x_g x_{g^{-1}h} h \\
&= \sum_{h \in G} \sum_{g \in \langle a \rangle} \chi_3(g) x_g x_{hg} h - \sum_{h \in G} \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{hg^{-1}} h \\
&\quad - \sum_{h \in G} \sum_{g \in \langle a \rangle} \chi_3(g) x_g x_{gh} h + \sum_{h \in G} \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h} h
\end{aligned}$$

as required. □

Lemma 36. *The coefficient of $h = a^k$ in $[\alpha_1, \alpha_4]$ is*

$$(-1)^{k+1} \sum_{g \in G \setminus \langle a \rangle} \chi_3(g^{-1}) x_g x_{g^{-1}h} + \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h}.$$

Proof. We compute

$$\begin{aligned}
&\sum_{g \in \langle a \rangle} \chi_3(g) x_g x_{hg} - \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{hg^{-1}} \\
&\quad - \sum_{g \in \langle a \rangle} \chi_3(g) x_g x_{gh} + \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h} \\
&= - \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{hg^{-1}} + \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h} \\
&= - \sum_{g \in G \setminus \langle a \rangle} \chi_3(g^{-1}h) x_{g^{-1}h} x_{hh^{-1}g} + \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h} \\
&= (-1)^{k+1} \sum_{g \in G \setminus \langle a \rangle} \chi_3(g^{-1}) x_g x_{g^{-1}h} + \sum_{g \in G \setminus \langle a \rangle} \chi_3(g) x_g x_{g^{-1}h}
\end{aligned}$$

as required. □

From Lemma 36, we know that

$$[\alpha_1, \alpha_4] \neq 0.$$

From Theorems 29 and 30, we think that degree one representations form natural pairing (χ_1, χ_2) and (χ_3, χ_4). Furthermore, we will have another interesting algebraic properties soon.

Lemma 37. *The coefficient of $h = a^k b$ in $[\alpha_1, \alpha_4]$ is*

(1) *For the case that m is odd.*

$$\begin{cases} -(1+i)\sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{gh} + (1+i)\sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{g^{-1}h}, & \text{if } k \text{ is odd.} \\ -(1-i)\sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{gh} + (1-i)\sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{g^{-1}h}, & \text{if } k \text{ is even.} \end{cases}$$

(2) *For the case that m is even.*

$$\begin{cases} 2\sum_{g\in G}\chi_3(g)x_g x_{g^{-1}h}, & \text{if } k \text{ is odd.} \\ 0, & \text{if } k \text{ is even.} \end{cases}$$

Proof. (1) For the case that m is odd, we compute

$$\begin{aligned} &\sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{hg} - \sum_{g\in G\setminus\langle a\rangle}\chi_3(g)x_g x_{hg^{-1}} \\ &\quad - \sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{gh} + \sum_{g\in G\setminus\langle a\rangle}\chi_3(g)x_g x_{g^{-1}h} \\ &= \sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{g^{-1}h} - \sum_{g\in\langle a\rangle}\chi_3(gb)x_{gb}x_{hb^{-1}g^{-1}} \\ &\quad - \sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{gh} + \sum_{g\in\langle a\rangle}\chi_3(gb)x_{gb}x_{b^{-1}g^{-1}h} \\ &= \sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{g^{-1}h} - i\sum_{g\in\langle a\rangle}\chi_3(g)x_{gb}x_{a^k g^{-1}} \\ &\quad - \sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{gh} + i\sum_{g\in\langle a\rangle}\chi_3(g)x_{gb}x_{a^{-k}g} \\ &= \sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{g^{-1}h} - i\sum_{g\in\langle a\rangle}\chi_3(ga^k)x_{ga^k b}x_{g^{-1}} \\ &\quad - \sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{gh} + (-1)^k i\sum_{g\in\langle a\rangle}\chi_3(g)x_{gh}x_g \\ &= \sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{g^{-1}h} - (-1)^k i\sum_{g\in\langle a\rangle}\chi_3(g)x_{gh}x_{g^{-1}} \\ &\quad - \sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{gh} + (-1)^k i\sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{gh} \end{aligned}$$

as required.

(2) For the case that m is even, we compute

$$\sum_{g\in\langle a\rangle}\chi_3(g)x_g x_{hg} - \sum_{g\in G\setminus\langle a\rangle}\chi_3(g)x_g x_{hg^{-1}}$$

$$
\begin{aligned}
&- \sum_{g\in\langle a\rangle} \chi_3(g) x_g x_{gh} + \sum_{g\in G\setminus\langle a\rangle} \chi_3(g) x_g x_{g^{-1}h} \\
&= \sum_{g\in\langle a\rangle} \chi_3(g) x_g x_{g^{-1}h} - \sum_{g\in\langle a\rangle} \chi_3(gb) x_{gb} x_{hb^{-1}g^{-1}} \\
&\quad - \sum_{g\in\langle a\rangle} \chi_3(gb) x_{b^{-1}bg} x_{gh} + \sum_{g\in G\setminus\langle a\rangle} \chi_3(g) x_g x_{g^{-1}h} \\
&= \sum_{g\in G} \chi_3(g) x_g x_{g^{-1}h} - \sum_{g\in\langle a\rangle} \chi_3(g) x_{gb} x_{a^k g^{-1}} \\
&\quad - \sum_{g\in\langle a\rangle} \chi_3(gb) x_{b^{-1}g^{-1}b} x_{ga^k b} \\
&= \sum_{g\in G} \chi_3(g) x_g x_{g^{-1}h} - \sum_{g\in\langle a\rangle} \chi_3(ga^k) x_{ga^k b} x_{g^{-1}} \\
&\quad - \sum_{g\in\langle a\rangle} \chi_3(ga^{-k}b) x_{b^{-1}g^{-1}a^k b} x_{gb} \\
&= \sum_{g\in G} \chi_3(g) x_g x_{g^{-1}h} + (-1)^{k+1} \sum_{g\in\langle a\rangle} \chi_3(g) x_g x_{g^{-1}h} \\
&\quad + (-1)^{k+1} \sum_{g\in G\setminus\langle a\rangle} \chi_3(g) x_{g^{-1}h} x_g \\
&= \sum_{g\in G} \chi_3(g) x_g x_{g^{-1}h} + (-1)^{k+1} \sum_{g\in G} \chi_3(g) x_g x_{g^{-1}h}
\end{aligned}
$$

as required. □

We have the following algebraic properties.

Theorem 38. *The formulas*

$$
\begin{aligned}
[\alpha_1, \alpha_3 + \alpha_4] &= 0, \\
[\alpha_2, \alpha_3 + \alpha_4] &= 0, \\
[\alpha_3, \alpha_1 + \alpha_2] &= 0, \\
[\alpha_4, \alpha_1 + \alpha_2] &= 0
\end{aligned}
$$

hold.

Proof. By Lemmas 33, 34, 36 and 37, we have

$$[\alpha_1, \alpha_3 + \alpha_4] = 0.$$

From Lemmas 25 and 26, this completes the proof. □

Acknowledgments I am deeply grateful to Prof. Hiroyuki Ochiai and Prof. Minoru Itoh who provided the helpful comments and suggestions. Also, I would like to thank my college in the Graduate School of Mathematics of Kyushu University, in particular Cid Reyes, Tomoyuki Tamura, Yuka Suzuki for comments and suggestions. I would also like to express my gratitude to my family for their moral support and warm encouragements.

REFERENCES

[1] Benjamin Steinberg, *Representation Theory of Finite Groups*. Springer, 2012.
[2] C. Shan, C. Hong, T. Guoping, Augmentation quotients for complex representation rings of dihedral groups, Frontiers of Mathematics in China. Vol. 7, pp. 1-18, 2012.
[3] K. W. Jonson, On the group determinant, Mathematical Proceedings of the Cambridge Philosophical Society. Vol. 109, pp. 299-311, 1991.
[4] Z. Qingxia, Y. Hong, On the Structure of Augmentation Quotient Groups for the Generalized Quaternion Group, Algebra Colloquium. Vol. 19, pp. 137-148, 2012.

Naoya Yamaguchi
Graduate School of Mathematics
Kyushu University
Nishi-ku, Fukuoka 819-0395
Japan
n-yamaguchi@math.kyushu-u.ac.jp